

A service of the U.S. National Institutes of Health
[Sign In](#) [Register](#)

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC
 Journals Books

Search for

[Advanced Search \(beta\)](#)

[Limits](#) [Preview/Index](#) [History](#) [Clipboard](#) [Details](#)

Note: Performing your original search, *intraperitoneal injection study*, in PubMed will retrieve 13537 citations.

Display Show Sort By Send to

All: 1 Review: 0

☐ 1: [Gan To Kagaku Ryoho](#). 1998 May;25(6):881-5.

[Links](#)

[Experimental study on 5-FU intraperitoneal injection for liver metastasis]

[Article in Japanese]

Kawahara H, Hirai K, Aoki T.

Dept. of Second Surgery, Jikei University School of Medicine.

The 5-FU concentrations in peripheral blood and portal blood were determined in rats after 5-FU injection via three routes. Rats were given 1 ml of 5-FU (250 mg/kg) via the subphrenic, Douglas or intravenous routes. From comparison of 5-FU concentrations between the intravenous injection group and the intraperitoneal group (subphrenic injection group + Douglas injection group), the intraperitoneal group had a lower concentration than the intravenous one in peripheral blood. The intraperitoneal group showed a higher concentration than the intravenous one in portal blood. The intraperitoneal group was divided into two groups, a subphrenic injection group and a Douglas injection group. Though there was no significant difference between the 2 groups in the 5-FU concentrations in portal blood, the 5-FU concentration of peripheral blood in subphrenic injection group was significantly higher than in the Douglas injection group. It was conceivable that the subphrenic injection group had another absorption route from the abdominal cavity to the peripheral blood. Accordingly, the intraperitoneal injection of the agent may be useful therapy for liver metastasis. However, due care must be given to the injection lesion in the abdominal cavity, the concentration and volume of the agent.

PMID: 9617327 [PubMed - indexed for MEDLINE]

Related Articles

[Experimental study on CDDP, 5-FU concentrations of paraaortic and mesenteric lymph nodes after intraperitoneal infusion] [\[Nippon Onkologu Gakkaishi. 1996\]](#)

Does the administration route of leucovorin have any influence on the impairment of colonic healing caused by intraperitoneal 5-fluorouracil treatment? [\[Eur Surg Res. 2001\]](#)

Effect of carcinomatosis and intraperitoneal 5-fluorouracil on peritoneal blood flow modulated by vasopressin in the rat as measured with laser Doppler flow technique [\[J Surg Res. 2004\]](#)

5-FU uptake in liver metastases after intravenous and intraperitoneal administration: an autoradiographic study in the rat. [\[Anticancer Res. 1996\]](#)

[Concentrations of 5-fluorouracil (5-FU) in serum and tissues at venous injection of tegafur or 5-FU--clinical study on colorectal cancer] [\[Gan To Kagaku Ryoho. 1998\]](#)

[» See all Related Articles...](#)

Patient Drug Information

» Fluorouracil (Adrucil®) Your doctor has ordered the drug fluorouracil to help treat your illness. The drug is given by injection into a vein.

[» read more ...](#)

Display Show Sort By Send to

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)